

# A predictive risk model for first line treatment failure in South Africa

CROI 2015  
Poster #1120

Julia Rohr<sup>1,5</sup>, Prudence Ive<sup>2</sup>, C. Robert Horsburgh<sup>1,5</sup>, Rebecca Berhanu<sup>2</sup>, Kate Shearer<sup>3</sup>, Mhairi Maskew<sup>3</sup>, Lawrence Long<sup>3</sup>, Ian Sanne<sup>1,2,3,4</sup>, Matthew P Fox<sup>1,3,5</sup>

<sup>1</sup>Center for Global Health & Development, Boston University, Boston, MA, USA; <sup>2</sup>Clinical HIV Research Unit, Department of Internal Medicine, School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa; <sup>3</sup>Health Economics and Epidemiology Research Office, Department of Internal Medicine, School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa; <sup>4</sup>Right to Care, Johannesburg, South Africa; <sup>5</sup>Department of Epidemiology, Boston University School of Public Health, Boston, MA, USA

## INTRODUCTION

### Background

Although individual predictors of first line antiretroviral therapy (ART) failure have been identified, few studies in resource-limited settings have been large enough for predictive modeling. Understanding the absolute risk of first line failure is useful for patient monitoring and for effectively targeting limited resources for second line ART.

### Objectives

- 1) To estimate absolute risk of failure of first line ART over 5 years on ART as a function of key demographic, clinical, and immunologic factors at the start of ART
- 2) To develop a predictive model for other South African clinics to identify patients at high risk of failure.

## METHODS

- **Design:** Observational cohort
- **Data:** Medical records from 9 South African public HIV clinics
- **Inclusion:**
  - Patients initiated on first line ART after 2004 with at least 6 months of follow-up time.
- **Accelerated failure time models**
- **Person time**
  - Start: 6 months after ART start
  - End: 2 consecutive viral loads >1000 copies/mL

### Candidate Predictor Variables

**Demographic:** Sex, age, history of smoking and alcohol use, pregnancy  
**Lab values:** CD4, hemoglobin, RBC, MCV, AST, ALT, creatinine clearance  
**Clinical measures:** WHO stage, body mass index (BMI), history of TB, peripheral neuropathy

**Treatment:** ARVs in first line regimen, missed visits in first 6 months on treatment

### Model Development

1. Diagnostic plots to choose survival distribution to fit data for accelerated failure time models
2. Unadjusted accelerated failure time models for individual predictors (stratified by clinic)
3. Adjusted accelerated failure time model (stepwise selection of predictors)
4. Impute missing data
5. Re-run adjusted models
6. Cross validation of models w/imputed data
  - > Develop model on all clinics but one
  - > Test model on excluded clinic
  - > Rotate excluded clinic (9 times)
  - > Output weighted average of:
    - Harrell's C statistic
    - % of patients for whom model correctly discriminates failure
    - Difference between actual and predicted 5-year failure

## RESULTS

### Final Model

Accelerated failure time model estimates used to calculate scores for risk groups:

### RISK SCORE CALCULATION


<b>1. Sex &amp; age:</b>		<b>3. CD4 count:</b>		
Females: Age:		0-25	10	
18-24.9	8	25-50	8	
25-29.9	6	51-100	6	
30-34.9	4	101-200	4	
35-39.9	2	201-350	2	
40-44.9	1	>350	0	
45+	0			
Males: Age:		<b>4. MCV:</b>		
18-24.9	7	<80	0	
25-29.9	6	80-95	1	
30-34.9	5	95+	3	
35-49.9	4			
50+	3			
<b>2. NNRTI:</b>		<b>5. Hemoglobin:</b>		
NVP	3	<12	1	
EFV	0	12+	0	
		<b>6. History of TB:</b>		
		Yes	1	
		No	0	
		<b>7. Missed visits during first 6 months on treatment:</b>		
		Yes	4	
		No	0	

SCORE	RISK GROUP
>17	High
14-17	Medium-high
12-13	Middle
9-11	Medium-low
0-8	Low

### Prediction of 5-Year Treatment Failure

Risk group:	Years on ART:				
	1	2	3	4	5
<b>High</b>	3.8%	10.1%	15.4%	20.1%	24.4%
<b>Medium-high</b>	2.7%	7.3%	11.2%	14.7%	18.0%
<b>Medium</b>	2.2%	5.9%	9.1%	12.1%	14.8%
<b>Medium-low</b>	1.8%	4.9%	7.5%	10.0%	12.3%
<b>Low</b>	1.4%	3.7%	5.7%	7.6%	9.4%

Comparison between model-predicted treatment failure and actual failure over time on ART for study population



## CONCLUSION

The predictive model was able to discriminate between patients at higher risk of first line virologic failure. 24% of patients at highest risk failed treatment in 5 years. Identification of patients at highest risk of failure is useful for patient monitoring, referral for adherence counseling, and targeting adherence interventions to improve patient outcomes and avoid need for second line ART.

Center for Global Health and Development, Boston University School of Public Health, 801 Massachusetts Ave 3<sup>rd</sup> Floor, Boston, MA 02118 USA, 617-414-1252; [jrohr@bu.edu](mailto:jrohr@bu.edu); <http://www.bu.edu/cghd>

Funding was provided by USAID under the terms of Cooperative Agreement 674-A-00-09-00018-00 to Boston University and Cooperative Agreement 674-A-00-02-00018 and CA 674 A 00 08 0000 700 to Right to Care; INROADS USAID-674-A-12-00029 and the National Institute of Allergy and Infectious Diseases (NIAID) Award Number K01AI08309. This study is made possible by the generous support of the American people through the United States Agency for International Development (USAID) and the National Institutes of Health. The contents are the responsibility of the authors and do not necessarily reflect the views of USAID, NIH or the United States government.

